

## WEST

☐  

L6: Entry 4 of 14

File: USPT

Jan 9, 2001

US-PAT-NO: 6171618

DOCUMENT-IDENTIFIER: US 6171618 B1

TITLE: Combination dosage form comprising cetirizine and pseudoephedrine

DATE-ISSUED: January 9, 2001

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Johnson; Barbara A.	Niantic	CT		
Korsmeyer; Richard W.	Old Lyme	CT		
Oksanen; Cynthia A.	Mystic	CT		

US-CL-CURRENT: 424/472; 424/480

## CLAIMS:

What is claimed is:

1. A solid dosage form comprising cetirizine and pseudoephedrine

wherein at least a portion of said pseudoephedrine is contained in a core comprising said portion, whereby release of said portion of pseudoephedrine into an environment of use is sustained;

wherein said cetirizine is contained as an immediate-release component in said dosage form; and

wherein said dosage form is substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof, said alcohol and/or derivatives being present in an amount less than that required to react with 5% of the cetirizine in the dosage form.

2. A dosage form as defined in claim 1, wherein said cetirizine is in the form of a pharmaceutically acceptable salt.

3. A dosage form as defined in claim 2, wherein said salt

is the dihydrochloride.

4. A dosage form as defined in claim 1, wherein said pseudoephedrine is in the form of a pharmaceutically acceptable salt.

5. A dosage form as defined in claim 4, wherein said salt is the sulfate or the hydrochloride.

6. A dosage form as defined in claim 1, wherein said dosage form is substantially free of alcohols having a molecular weight lower than 250 and reactive derivatives thereof.

7. A dosage form as defined in claim 6, wherein said dosage form is substantially free of alcohols having a molecular weight lower than 500 and reactive derivatives thereof.

8. A dosage form as defined in claim 7, wherein said dosage form is substantially free of alcohols having a molecular weight lower than 1000 and reactive derivatives thereof.

9. A dosage form as defined in claim 1, wherein all of said pseudoephedrine is contained in said core.

10. A dosage form as defined in claim 1, wherein said core comprises a pseudoephedrine-containing immediate release composition surrounded by a rate limiting membrane.

11. A dosage form as defined in claim 1, wherein said core comprises a pseudoephedrine-containing sustained release matrix.

12. A dosage form as defined in claim 11, wherein said matrix comprises a hydrophilic polymer.

13. A dosage form as defined in claim 1, in the form of a tablet.

14. A dosage form as defined in claim 1, in the form of a multiparticulate.

15. A dosage form as defined in claim 1, wherein said

portion of pseudoephedrine is released into said environment of use over a period of 4 to 36 hours.

16. A dosage form as defined in claim 15, wherein said period is about 8 to about 24 hours.

17. A solid dosage form comprising cetirizine and pseudoephedrine, wherein at least a portion of said pseudoephedrine is contained in a core surrounded by a permeable membrane, whereby release of said portion of pseudoephedrine into an environment of use is sustained;

wherein said cetirizine is contained as an immediate-release component in said dosage form; and

wherein said dosage form is substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof, said alcohol and/or derivatives being present in an amount less than that required to react with 5% of the cetirizine in the dosage form.

18. A dosage form as defined in claim 17, wherein said cetirizine is in the form of a pharmaceutically acceptable salt.

19. A dosage form as defined in claim 18, wherein said salt is the dihydrochloride.

20. A dosage form as defined in claim 17, wherein said pseudoephedrine is in the form of a pharmaceutically acceptable salt.

21. A dosage form as defined in claim 20, wherein said salt is the sulfate or the hydrochloride.

22. A dosage form as defined in claim 17, wherein said dosage form is substantially free of alcohols having a molecular weight lower than 250 and reactive derivatives thereof.

23. A dosage form as defined in claim 22, wherein said dosage form is substantially free of alcohols having a molecular weight lower than 500 and reactive derivatives thereof.

24. A dosage form as defined in claim 23, wherein said dosage form is substantially free of alcohols having a molecular weight lower than 1000 and reactive derivatives thereof.

25. A dosage form as defined in claim 17, in the form of a tablet.

26. A dosage form as defined in claim 25, wherein said core comprises about 60 to about 360 mg of pseudoephedrine, about 70 to about 425 mg of compressible excipients, about 2 to about 17 mg of binder, and about 0.5 to about 4 mg of lubricant.

27. A dosage form as defined in claim 17, in the form of a multiparticulate.

28. A dosage form as defined in claim 17, wherein said permeable membrane is fabricated from a cellulosic material.

29. A dosage form as defined in claim 17, wherein said portion of pseudoephedrine is released into said environment of use over a period of 4 to 36 hours.

30. A dosage form as defined in claim 23, wherein said period is about 8 to about 24 hours.

31. A process for making a solid dosage form comprising cetirizine and pseudoephedrine, comprising coating a sustained release core comprising pseudoephedrine, with an immediate release layer comprising cetirizine and a water soluble film forming polymer, said layer and said core being substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof, said alcohol and/or derivatives being present in an amount less than that required to react with 5% of the cetirizine in the dosage form.

32. A process as defined in claim 31, wherein said dosage form is a tablet.

33. A process as defined in claim 31, wherein said dosage form is a multiparticulate.

34. A process for making a solid dosage form containing

cetirizine and pseudoephedrine, comprising the steps of coating a shaped composition comprising pseudoephedrine with a permeable membrane,

coating said permeable membrane with an immediate release layer comprising cetirizine and a water soluble film forming polymer,

said composition, said membrane and said layer being substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof.

35. A process as defined in claim 34, wherein said dosage form is a tablet.

36. A process as defined in claim 35, wherein said core comprises about 60 to about 360 mg of pseudoephedrine, about 70 to about 425 mg of compressible excipients, about 2 to about 17 mg of binder, and about 0.5 to about 4 mg of lubricant.

37. A process as defined in claim 34, wherein said dosage form is a multiparticulate.

38. A process as defined in claim 34, wherein said permeable membrane is fabricated from a cellulosic material.

39. A dosage form as defined in claim 1, containing an amount of said alcohol and/or derivatives less than the amount required to react with 1% of the cetirizine in the dosage form.

40. A dosage form as defined in claim 39, containing an amount of said alcohol and/or derivatives less than the amount required to react with 0.5% of the cetirizine in the dosage form.

41. A dosage form as defined in claim 40, which is totally free of said alcohol and/or derivatives.

42. A dosage form as defined in claim 17, containing an amount of said alcohol and/or derivatives less than the amount required to react with 1% of the cetirizine in the dosage form.

43. A dosage form as defined in claim 42, containing an amount of said alcohol and/or derivatives less than the amount required to react with 0.5% of the cetirizine in the dosage form.

44. A dosage form as defined in claim 43, which is totally free of said alcohol and/or derivatives.

45. A process as defined in claim 31, wherein said dosage form contains an amount of said alcohol and/or derivatives less than the amount required to react with 1% of the cetirizine therein.

46. A process as defined in claim 45, wherein said dosage form contains an amount of said alcohol and/or derivatives less than the amount required to react with 0.5% of the cetirizine therein.

47. A process as defined in claim 46, wherein said dosage form is totally free of said alcohol and/or derivatives.

## WEST



Generate Collection

Print

L6: Entry 4 of 14

File: USPT

Jan 9, 2001

DOCUMENT-IDENTIFIER: US 6171618 B1

TITLE: Combination dosage form comprising cetirizine and pseudoephedrineAbstract Text (1):

A dosage form containing cetirizine as an immediate release component and pseudoephedrine or a pharmaceutically acceptable salt thereof as a controlled release component. A portion of the pseudoephedrine can also be incorporated as an immediate release component. The dosage form is free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof.

Brief Summary Text (2):

This invention relates to dosage forms comprising cetirizine and pseudoephedrine, containing both a sustained release and an immediate release component.

Brief Summary Text (6):

Pseudoephedrine, as well as pharmaceutically acceptable acid addition salts thereof such as the hydrochloride and sulfate salts, is a sympathomimetic drug known by those skilled in the art as a safe therapeutic agent for treating nasal congestion. It is commonly administered orally and concomitantly with an antihistamine for treatment of nasal congestion for the treatment of allergic rhinitis.

Brief Summary Text (7):

Cetirizine and pseudoephedrine can be administered together. The general schedule for administering the two drugs together involves one 10 mg cetirizine tablet per day, plus eight 30 mg immediate-release pseudoephedrine tablets divided into four doses over the course of a day. It is well known, however, that patient compliance with a multiple dose daily administration schedule can be affected by the inconvenience of having to remember to take medicine at numerous, appropriately spaced intervals. It would accordingly be useful if patients could take cetirizine plus pseudoephedrine as a unitary dosage form, such as a tablet, say once-daily or twice-daily, to improve convenience and better ensure patient compliance.

Brief Summary Text (9):

In one aspect, this invention provides a solid dosage form comprising cetirizine and pseudoephedrine

Brief Summary Text (10):

wherein at least a portion of said pseudoephedrine is contained in a core comprising said portion of pseudoephedrine, whereby release of said pseudoephedrine into an environment of use is sustained;

Brief Summary Text (11):

wherein said cetirizine is contained as an immediate-release component in said dosage form; and

Brief Summary Text (13):

In a preferred embodiment, this invention provides a solid dosage form comprising cetirizine and pseudoephedrine,

Brief Summary Text (14):

wherein at least a portion of said pseudoephedrine is contained in a core comprising said portion of pseudoephedrine, said core being surrounded by a permeable membrane, whereby release of said pseudoephedrine into an environment of use is sustained;

Brief Summary Text (15):

wherein said cetirizine is contained as an immediate-release component in said dosage form; and

Brief Summary Text (18):

This invention further provides a process for making a solid dosage form containing cetirizine and pseudoephedrine, comprising

Brief Summary Text (19):

coating a shaped sustained release core comprising pseudoephedrine with an immediate release layer comprising cetirizine and a water soluble film forming polymer, said core and said immediate-release layer being substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof.

Brief Summary Text (20):

In a preferred embodiment this invention provides a preferred process for making a solid dosage form containing cetirizine and pseudoephedrine, comprising the steps of

Brief Summary Text (21):

coating a shaped composition comprising pseudoephedrine with a permeable membrane

Brief Summary Text (22):

coating said permeable membrane with an immediate release layer comprising cetirizine and a water soluble film forming polymer,

Brief Summary Text (24):

"Cetirizine" as employed herein is intended to include not only the free compound of formula (I) but also any pharmaceutically acceptable salt thereof. Preferred are acid addition salts, especially the dihydrochloride (referred to also herein as the "hydrochloride"). "Cetirizine" is also intended to cover individual enantiomers as well as the racemate.

Brief Summary Text (25):

"Pseudoephedrine" as employed herein is intended to be inclusive of the free base and also of any pharmaceutically acceptable acid addition salt. Preferred for use in this invention are the sulfate and hydrochloride salts.

Brief Summary Text (26):

The term "alcohols" is used generically to mean any compound having one or more hydroxyl (--OH) groups reactive toward cetirizine. The term is thus inclusive of diols and polyols as well as mono-alcohols which are reactive toward cetirizine under normal conditions of processing and storage.

Brief Summary Text (27):

A "reactive derivative" as used herein includes those materials having alcohol (--OH) groups which have been esterified and which can react with cetirizine by transesterification.

Brief Summary Text (29):

The invention provides a dosage form containing both cetirizine and pseudoephedrine. The cetirizine is contained in an immediate-release component from which it starts to be delivered or released substantially immediately upon ingestion (i.e., upon swallowing). No mechanism has been incorporated into the dosage form to delay the release which would otherwise take place upon exposure to a use environment, such as the luminal fluid of the gastrointestinal (GI) tract. Generally the cetirizine should be at least 80% released from the dosage form within an hour after administration.

Brief Summary Text (30):

The pseudoephedrine, by contrast, releases in a sustained fashion, at least about 75% of the drug contained in the dosage form releasing over a period of 4 to 36 hours, preferably about 8 to about 24 hours, although the period of sustained release can be tailored to have an immediate release component, as disclosed further below. The term "about" as used above and elsewhere herein means plus or minus 10% for each of the numerical limits. The sustained release pseudoephedrine is contained in a core which can be engineered in a variety of ways and embodiments to implement sustained release. For example, the pseudoephedrine can be incorporated into a sustained release matrix that meters pseudoephedrine out over a period of 4 to 36 hours, the matrix thus constituting the core. Alternatively, the pseudoephedrine core can comprise a shaped pseudoephedrine immediate release composition and a surrounding, rate limiting membrane which imparts sustained release behavior to the core.

Brief Summary Text (31):

In the final product dosage form (i.e., the dosage form intended to be sold or administered), and during processing when making the dosage form, it is important to avoid contact of cetirizine with alcohols having a molecular weight less than 100 since such contact can result in a reaction with cetirizine, usually esterification, and thereby damage the dosage form by reacting with the active medicinal agent. Thus, the dosage form should be substantially free of such reactive components at the time the immediate-release cetirizine component is introduced into the dosage form, and thereafter. Alcohols and other



reactive components can be employed during processing, so long as they are removed to reduce or eliminate their presence prior to introducing cetirizine. Not employing alcohols at all during processing, or removing them prior to introducing cetirizine into the dosage form, is what is meant by the phrase "substantially free of alcohols having a molecular weight less than . . .". Some level of alcohols which are reactive with cetirizine can be tolerated in the final dosage form, the exact level depending on the particular alcohol. In general, to be "substantially free" of alcohol(s) the total amount of alcohol of molecular weight less than 100, whether a single alcohol or a mixture, should be less than the amount which would be required to react with 5% of the cetirizine in the dosage form, preferably less than the amount which would be required to react with 1% of the cetirizine. More preferably, it is preferred that the level of alcohol having a molecular weight less than 100, whether a single alcohol or a mixture, be less than that amount of alcohol required to react with 0.5% of the cetirizine in the dosage form, and most preferably that the dosage form be totally free of alcohols having a molecular weight less than 100.

Brief Summary Text (32):

Thus the final dosage form, in both the core and outside layer(s), must be substantially free of low molecular weight alcohols and reactive derivatives thereof. Such materials include relatively low molecular weight monohydric and polyhydric alcohols which are conventionally known and frequently, if not universally, used as solvents in the formulations arts, and compositions containing them as vehicles or carriers. Examples of such reactive alcohols include lower molecular weight alcohols such as methanol, ethanol, isopropanol, and glycerin. Because they are difficult to remove, high boiling point alcohols such as glycerin can be problematic, and it is preferred such components be avoided altogether. Many plasticizers are alcohols, and contact with cetirizine should accordingly be avoided as well if they have a low molecular weight. Many plasticizers are also esters, i.e., materials which are reactive derivatives of alcohols, and contact of cetirizine with these materials should be avoided as well since the ester groups can undergo transesterification with cetirizine and thereby damage the dosage form. Not all alcohol group-containing components are reactive toward cetirizine under normal conditions of processing and storage (temperatures less than 100.degree. C.), however, and such components are not "alcohols" for purposes of this invention. Examples include cellulosic materials containing free hydroxyl groups, such as microcrystalline celluloses used as compressible excipients as well as cellulose ethers and esters useful as coatings in the formulations described herein.

Brief Summary Text (34):

The cetirizine can be incorporated into the dosage form as an "Immediate release" component in a variety of ways. For example, it can be incorporated into an exterior coating for a tablet from which it releases substantially immediately upon ingestion. The coating can be all-covering, or can cover less than the surface of the dosage form, as described below. Such a coating can similarly be applied to each of the particles comprising a multiparticulate. If the dosage form is to be a capsule, the cetirizine can be contained in a single pellet inside the capsule from which it releases substantially immediately once the capsule shell dissolves. Alternatively, the cetirizine can be contained in several smaller pellets or be present as immediate release particles. In this type of capsule embodiment, the pseudoephedrine is generally present, for example, as a slow release multiparticulate, each particle comprising a central core of pseudoephedrine incorporated into a matrix or surrounded by a rate-limiting membrane.

Brief Summary Text (35):

The term "tablet" refers to the conventional macroscopic dosage form as known in the art, and as applied to this invention indicates a unitary dosage having a central core which releases pseudoephedrine in a sustained fashion, over a period of 4 to 36 hours, preferably about 8 to about 24 hours. As previously mentioned the core can be a matrix which meters out pseudoephedrine. Alternatively, the pseudoephedrine core can be formed from an immediate release, pseudoephedrine-containing composition which is surrounded by a water insoluble, permeable, rate-limiting membrane that provides for sustained release of pseudoephedrine by limiting the rate at which pseudoephedrine diffuses into the environment of use. The core is in turn coated over at least a portion of its surface with a layer comprising cetirizine and a water soluble film forming polymer that provides immediate release. The immediate release cetirizine-containing layer is preferably coated over the entire surface of the core for convenience in formulating the tablet. The immediate release layer can also, if desired, be made to cover less than the entire area of the membrane by conventional procedures known to the art. For example, if the tablet has flat surfaces, the cetirizine-containing layer can be coated exclusively onto one or more, but less than all, of the surfaces. If the tablet is spherical, the cetirizine-containing layer can be coated over less than the entire surface of the sphere. A tablet dosage form is desirable as a pre-made, readily available dosage form which can be administered in the majority of cases to patients who need and/or desire a medication containing both an antihistamine plus a decongestant.

Brief Summary Text (36):

In the case of a pseudoephedrine-containing core surrounded by a rate limiting membrane, the membrane is "permeable". Permeability can be achieved by implementing a coating which completely surrounds the core and which has a pore size that allows the passage of both water and pseudoephedrine through the membrane. In this type of membrane, water crosses the membrane into the core and dissolves the pseudoephedrine so that pseudoephedrine is released back across the membrane into the GI tract at a desired rate, i.e, a rate which is reasonably constant until substantially all of the pseudoephedrine has been released over a period of 4 to 36 hours. This type of membrane is disclosed and described, for example, in EP-A-0 357

369, herein incorporated by reference in its entirety. Alternatively, the membrane can be permeable to water only, but have one or more openings such as a round hole or other shape opening, shape not being critical. The opening can be physically implemented, for example by drilling, somewhere on the surface to allow the release of pseudoephedrine. This type of dosage is known to the art and described in numerous U.S. patents including U.S. Pat. Nos. 4,915,954; 4,915,953; 4,915,952; 4,847,093; is and 4,810,502; each of which is herein incorporated by reference in its entirety.

Brief Summary Text (37):

The term "multiparticulate dosage form" or multiparticulate is intended to embrace a dosage form comprising a multiplicity of particles whose totality represents the intended therapeutically useful dose of cetirizine plus pseudoephedrine. The particles generally are of a diameter from about 50 microns to about 0.3 cm, with a preferred range of 100 .mu.m to 1 mm. Each individual particle is in essence a mini-tablet, and each comprises a central sustained release core of pseudoephedrine, with sustained release being implemented as described above. Each can also be coated, fully or partially, with an immediate release layer comprising cetirizine plus a water soluble film forming polymer.

Brief Summary Text (38):

In an alternative embodiment, a pseudoephedrine-containing matrix or membrane-coated multiparticulate, with no cetirizine-containing immediate-release coating, can be used to partially fill a capsule and one or more pellets of immediate-release cetirizine can be added as part of the capsule fill. A multiparticulate dosage form is desirable since it allows customizing or tailoring according to the weight of the patient by simply scaling the number of particles in the dosage form. Thus, for example, a weight of multiparticulate adapted for a particular patient or group of patients who may need more or less of the medication than that supplied in a pre-made tablet, can be used as fill for a capsule. Capsules adapted for larger or smaller patients can be tailored by simply adjusting, up or down as appropriate, the fill weight of multiparticulate.

Brief Summary Text (39):

If a multiparticulate is membrane-coated the individual particles are typically formed so that the membrane completely surrounds the central pseudoephedrine core and, in the use environment, is permeable to both water and pseudoephedrine.

Brief Summary Text (40):

In one dosage form embodiment of the invention, all of the cetirizine is incorporated into a separate (i.e, from the sustained release core) immediate-release coating that surrounds the pseudoephedrine core of the dosage form, and all of the pseudoephedrine is incorporated into the core. This embodiment can take the form of a macroscopic tablet, or the form of a multiparticulate for use as capsule fill.

Brief Summary Text (41):

In an alternative embodiment, all of the cetirizine and part of the pseudoephedrine is incorporated into the separate layer immediate release coating and the remainder of the pseudoephedrine is incorporated into the sustained release core. This alternative embodiment provides some of the pseudoephedrine for immediate release along with the immediate-release cetirizine also in the coating.

Brief Summary Text (42):

The amount of cetirizine administered can vary with the size of the patient, as discussed above, but will generally be in the range of about 5 to about 20 mg/day. The amount of pseudoephedrine administered will generally vary from about 60 to about 240 mg/day. Up to about 25% of the pseudoephedrine, corresponding to 15-60 mg/day, can be incorporated into the immediate release cetirizine-containing layer, thereby providing that amount for immediate release together with the cetirizine. It is preferred to incorporate all of the pseudoephedrine into the core, however.

Brief Summary Text (43):

The fact that alcohols and derivatives thereof such as esters are deleterious to formulations containing cetirizine is surprising in view of the fact that esters, for example glycerol esters, of cetirizine are difficult to make by conventional direct esterification methods. In view of the difficulty associated with direct esterification, it is surprising that cetirizine reactivity towards alcohols is such that extensive ester formation was observed when using processes to make dry solid formulations, such as tablets, in which cetirizine was exposed to glycerin or other alcohols in one or more steps.

Brief Summary Text (45):

Any dosage form useful in this invention can be tested to determine if a particular alcohol is potentially deleterious by the following procedure. The dosage form can be stored at 50.degree. C. for twelve weeks in a closed container (including the commercial container, if closable) and then dissolved in water. Following filtration, an aqueous sample can be injected in a liquid chromatograph and analyzed for the presence of cetirizine esters by reverse phase high performance liquid chromatography (HPLC) using UV detection at 231 nm. For the reverse phase system, a column having a C-18 hydrocarbon stationary phase, of the type designated as an Ultrasphere (registered trademark of Beckman Instruments, Inc.) C-18 ODS column, can be employed using an isocratic mobile phase consisting of 60% water having a pH of 2.5 (pH adjusted by adding

potassium dihydrogen phosphate) and 40% acetonitrile. Flow rate is typically 1 mL/minute. Cetirizine esters usually elute after cetirizine. Identification can be by comparison of an unknown peak with a known standard which elutes at the same retention time. Confirmation can be by mass spectrometry.

Brief Summary Text (46):

Sustained release cores are made from a matrix material in which pseudoephedrine is embedded or dispersed and which acts to retard the release of pseudoephedrine into the environment of use, the GI tract. When pseudoephedrine is embedded in a matrix of this sort, release of the drug takes place principally from the surface of the matrix. Thus the drug is released from the surface of the matrix after it diffuses through or when the surface of the matrix erodes, exposing the drug. In some matrices, both mechanisms can operate simultaneously. These types of systems are well known to the art and are discussed extensively, for example, in PCT/IB95/00264, published as WO 95/30422, herein incorporated by reference in its entirety.

Brief Summary Text (47):

A preferred embodiment of a matrix core has the form of a hydrophilic matrix, i.e., a core containing pseudoephedrine and, as matrix, an amount of hydrophilic polymer sufficient to provide a useful degree of control over the pseudoephedrine dissolution. Hydrophilic polymers useful for forming the matrix include hydroxypropyl methylcellulose (HPMC), hydroxypropyl cellulose (HPC), poly(ethylene oxide), poly(vinyl alcohol), xanthan gum, carbomer, carrageenan, and zooglan. A preferred material is HPMC. Other similar hydrophilic polymers may also be employed. In use, the hydrophilic material is swollen by, and eventually dissolves in, water. The pseudoephedrine is released both by diffusion from the matrix and by erosion of the matrix. The pseudoephedrine dissolution rate from a hydrophilic matrix core may be controlled by the amount and molecular weight of hydrophilic polymer employed. In general, using a greater amount of the hydrophilic polymer decreases the dissolution rate, as does using a higher molecular weight polymer. Using a lower molecular weight polymer increases the dissolution rate. The dissolution rate may also be controlled by the use of water soluble additives such as sugars, salts, or soluble polymers. Examples of these additives are sugars such as lactose, sucrose, or mannitol, salts such as NaCl, KCl, NaHCO<sub>3</sub>, and water soluble polymers such as poly(N-vinyl-2-pyrrolidone) or polyvinylpyrrolidone, low molecular weight HPC or HMPC or ethylcellulose. In general, increasing the fraction of soluble material in the formulation increases the release rate. A hydrophilic matrix core typically comprises about 20 to 90% by weight of pseudoephedrine and about 80 to 10% by weight of polymer.

Brief Summary Text (48):

A pseudoephedrine hydrogel matrix core can be coated with an immediate release coating comprising cetirizine and a water soluble film forming polymer.

Brief Summary Text (49):

A preferred matrix composition for use as a core comprises, by weight, about 50% to about 80% pseudoephedrine, about 15% to about 35% HPMC, 0% to about 35% lactose, 0% to about 15% PVP, 0% to about 20% microcrystalline cellulose, and about 0.25% to about 2% magnesium stearate.

Brief Summary Text (51):

In a particular embodiment, a pseudoephedrine matrix core can be coated with an impermeable coating, and an orifice (for example, a circular hole or a rectangular opening) can be provided by which the content of the tablet is exposed to the aqueous GI tract. This embodiment is along the lines of those presented in U.S. Pat. No. 4,792,448 to Ranade, herein incorporated by reference. The opening is typically of a size such that the area of the exposed underlying pseudoephedrine composition constitutes less than about 40% of the surface area of the device, preferably less than about 15%. The entire core can be coated with an immediate release layer comprising cetirizine and a water soluble film forming polymer.

Brief Summary Text (52):

In a further particular embodiment, a pseudoephedrine hydrogel matrix core can be coated with an impermeable material and an opening for drug transport produced by drilling a hole through the coating. The hole may be through the coating only, or may extend as a passageway into the core. The entire core can be coated with an immediate release layer comprising cetirizine and a water soluble film forming polymer.

Brief Summary Text (53):

In a further particular embodiment, a pseudoephedrine hydrophilic matrix core can be coated with an Impermeable material and a passageway for drug transport produced by drilling a passageway through the entire tablet. The entire core can be coated with an immediate release layer comprising cetirizine and a water soluble film forming polymer.

Brief Summary Text (54):

In a further particular embodiment, a pseudoephedrine hydrogel matrix core can be coated with an impermeable material and one or more passageways for drug transport produced by removing one or more strips from the impermeable coating or by cutting one or more slits through the coating, preferably on the radial surface or land of the tablet. The entire core can be

coated with an immediate release layer comprising cetirizine and a water soluble film forming polymer.

Brief Summary Text (55):

In a further particular embodiment, a pseudoephedrine hydrogel matrix core can be shaped in the form of a cone and completely coated with an impermeable material. A passageway for drug transport can then be produced by cutting off the tip of the cone and the entire cone can be coated with an immediate release layer comprising cetirizine and a water soluble film forming polymer.

Brief Summary Text (56):

In a further particular embodiment, a pseudoephedrine hydrogel matrix core can be shaped in the form of a hemisphere, completely coated with an impermeable material, and a passageway for drug transport produced by drilling a hole in the center of the flat face of the hemisphere. The entire core can be coated with an immediate release layer comprising cetirizine and a water soluble film forming polymer.

Brief Summary Text (57):

In a further particular embodiment, a pseudoephedrine hydrogel matrix core can be shaped in the form of a half-cylinder and completely coated with an impermeable material. A passageway for drug transport can be produced by cutting a slit through or removing a strip from the impermeable coating along the axis of the half-cylinder or along the centerline of the flat face of the half-cylinder. The entire core can be coated with an immediate release layer comprising cetirizine and a water insoluble film forming polymer.

Brief Summary Text (59):

The term "impermeable material" means a material having sufficient thickness and impermeability to pseudoephedrine such that no significant transport of pseudoephedrine can take place through the material during the time scale of the intended drug release (i.e., 41-36 hours). Such a coating can be obtained by selecting a coating material with a sufficiently low diffusion coefficient for pseudoephedrine and applying it to sufficient thickness. Materials for forming the impermeable coating of these embodiments include substantially all materials in which the diffusion coefficient of pseudoephedrine is less than about  $10 \times 10^{-7}$  cm<sup>2</sup>/s. It is noted that the preceding diffusion coefficient can be amply sufficient for a matrix device, as discussed above. In a device of the type now under discussion which has been provided with a macroscopic opening, however, a material with this diffusion coefficient, and almost any membrane material that is not a liquid, looks to the contained pseudoephedrine, by contrast, as though it is impermeable because the majority of transport is through the opening. Preferred coating materials include insoluble film forming polymers and waxes. Especially preferred are thermoplastic polymers, such as poly(ethylene-co-vinyl acetate), poly(vinyl chloride), ethylcellulose, and cellulose acetate. These materials exhibit the desired low permeation rate toward pseudoephedrine when applied as coatings of thicknesses greater than about 100  $\mu$ m.

Brief Summary Text (60):

When making cores which comprise an immediate release pseudoephedrine composition surrounded by a rate limiting membrane, the types of membranes which can be employed are widely known in the art, for example from the patents and publications previously mentioned including EP 0 357 369, U.S. Pat. No. 4,847,093, and U.S. Pat. No. 4,915,953. The membranes can be fabricated from water insoluble film forming polymers, for example from olefin and vinyl type polymers, organosilicon polymers, polysulfones, polyamides, polyurethanes, cellulose esters, cellulose ethers and the like. The polymers can be condensation polymers or addition polymers. Specific non-cellulosic polymers include poly(methylmethacrylate), poly(butylmethacrylate), polyethylene, ethylene vinylacetate copolymer, poly(dimethylsiloxane), polypropylene, polyvinyl chloride, polyvinyl alcohol, ethylene vinylalcohol, and the like.

Brief Summary Text (63):

Whether in the form of a tablet or a multiparticulate, the type of tablet core now discussed constitutes a membrane-moderated or reservoir system. In this type of sustained release system a reservoir of pseudoephedrine is surrounded by a rate-limiting membrane which allows passage of pseudoephedrine and water. The pseudoephedrine traverses the membrane by mass transport mechanisms well known in the art, including but not limited to dissolution in the membrane followed by diffusion across the membrane. As previously stated for this invention, these individual reservoir system dosage forms may be large, as in the case of a tablet containing a single large reservoir, or multiparticulate, as in the case of a capsule containing a plurality of reservoir particles, each individually coated with a membrane. The coating can be non-porous, yet permeable to pseudoephedrine (for example pseudoephedrine may diffuse directly through the membrane), or it may be porous. The particular mechanism of transport which operates to achieve sustained release is not believed to be critical.

Brief Summary Text (64):

Tablets having membrane-moderated cores can also be made as osmotic delivery systems, that is, systems such as those previously discussed in which rate-limiting membrane surrounding the pseudoephedrine core is impermeable or poorly permeable to pseudoephedrine but easily permeable to water. Water can then diffuse across the membrane and dissolve the

pseudoephedrine in the core which then passes out of the tablet through one or more holes which have been drilled through or otherwise implemented through the surface of the tablet so that the hole exposes the interior of the pseudoephedrine core. The size and shape of the hole is not critical and is usually on the order of 0.05 mm to 2 mm. The term "hole" is intended to cover any exit passageway regardless of what it's termed (e.g., hole, orifice, bore, slit and so forth).

**Brief Summary Text (66):**

A particularly useful process for tailoring the porosity of the coating comprises adding a pre-determined amount of a finely-divided water soluble material, such as sugars, salts or water soluble polymers to a solution or dispersion (e.g., an aqueous latex) of the membrane-forming polymer to be used. Additional useful pore formers include dimethyl sulfone, and nicotinamide. When the dosage form is ingested into the aqueous medium of the GI tract, these water soluble membrane additives are leached out of the membrane, leaving pores which facilitate release of the drug. The membrane coating can also be modified by the addition of plasticizer, as known in the art. These types of procedures are fully disclosed in U.S. Pat. Nos. 4,851,228, 4,687,660, and 3,538,214, all of which are herein incorporated by reference. Although alcohols are also known in the art to be useful as pore formers, including sorbitol, pentaerythritol, mannitol, and other aromatic and aliphatic mono-ols and polyols, including diols and triols, these types of alcohols as pore formers should be avoided in this process since they tend to react with cetirizine by esterification.

**Brief Summary Text (70):**

In the present invention, the rate-limiting membrane is to be an asymmetric membrane of the type described in EP 0 357 369, it is possible to pre-form the pores (as opposed to forming pores in-situ in the GI tract) by employing one or more, usually volatile, alcohols as pore-forming ingredients, but it is important that all alcohol be removed prior to applying the immediate-release layer which contains cetirizine. Thus if one or a mixture of low molecular weight alcohols is used as a pore former when making membrane-surrounded cores, the alcohol(s) should be removed after pore formation has been completed. Removal can be effected conventionally, for example by placing the cores in a forced air oven, vacuum oven, or fluidized bed for several hours, typically overnight. The alcohols employed should also be low boiling point, volatile liquids at room temperature such as methanol and ethanol. Higher boiling point alcohols which are liquid at room temperature, such as glycerol, butanol and higher mono-alcohols, and diols such as low molecular weight glycols like ethylene glycol and propylene glycol, are problematic and should be avoided altogether.

**Brief Summary Text (71):**

A tablet core according to the invention comprises about 60 to about 360 mg of pseudoephedrine, about 70 to about 425 mg of compressible excipients, about 2 to about 17 mg of binder, and about 0.5 to about 4 mg of lubricant. A preferred dosage form contains 240 mg of pseudoephedrine (or 120 mg if the dose is to be divided) as the hydrochloride and an amount of compressible excipient, binder, and lubricant within the ranges set above. Generally, the greater the amount of pseudoephedrine hydrochloride used in the core, the greater should be the corresponding amount of compressible excipient that is used.

**Brief Summary Text (76):**

If the immediate-release cetirizine component is implemented as a coating over the rate-limiting membrane, the coating can contain about 40 to about 85% by weight of cetirizine based on the weight of the coating, with the remainder (to 60 weight %) being a water soluble film forming polymer such as hydroxypropyl methylcellulose (HPMC) or a blend of hydroxypropyl cellulose and hydroxypropyl methylcellulose, which can be made or purchased under the registered trademark OPADRY, from Colorcon.

**Brief Summary Text (77):**

The water soluble film forming ingredient can also be a sugar. The coating can optionally contain other ingredients such as titanium dioxide or other suitable non-hydroxyl group containing pigment. For a (macroscopic) tablet, this coating will typically comprise from 2.5 to mg of cetirizine and 4 to 50 mg of film forming polymer.

**Brief Summary Text (78):**

In a preferred wet granulation process for making tablets, tablet cores are first formed by blending pseudoephedrine as the hydrochloride salt with microcrystalline cellulose (e.g., Avicel.RTM. PH101) and a binder (e.g., Klucel.RTM. EXF, registered trademark of Aqualon Company) in a V-blender or a P/K processor (Patterson-Kelley). The resulting blend is then granulated with water (approximately 9-13% by weight) to make a damp mass and dried to a final moisture content of less than 1% by weight as determined from a moisture balance. The granules are then milled and blended with additional microcrystalline cellulose such as Avicel.RTM. PH 200. Lubricant can then be added and the entire mass blended in a V-blender, and non-membrane coated cores then formed by compressing the resulting blend on a tablet press.

**Brief Summary Text (81):**

A solution of water soluble film forming agent (Opadry.RTM.) and cetirizine in water can then be sprayed onto the dried membrane-coated cores to apply the cetirizine coating. At this point the tablets are ready for therapeutic use. If desired,

however, an additional outermost taste masking layer can be added to the tablet or to the multiparticulate. Such layer can be added in a facile manner by simply coating an additional amount of film forming polymer as an all-covering layer over the cetirizine-containing layer. If desired opacifiers such as titanium dioxide can be added to the outermost coating.

**Brief Summary Text (82):**

Multiparticulates can be made per the description given above for tablets, wherein each multiparticulate is in essence a mini-tablet matrix core or a core surrounded by a membrane. A preferred process for manufacturing multiparticulate cores is the extrusion/spheronization process. For this process, pseudoephedrine is wet-massed with a binder, extruded through a perforated plate or die, and placed on a rotating disk. The extrudate ideally breaks into pieces which are rounded into spheres, spheroids, or rounded rods on the rotating plate. An example of this type of process and a composition involves using water to wet granulate a blend comprising, as a matrix, about to 75% of microcrystalline cellulose blended with, correspondingly, about 90 to 25% pseudoephedrine as the hydrochloride salt.

**Brief Summary Text (83):**

A further preferred process for manufacturing matrix multiparticulates is the preparation of wax granules. In this process, a desired amount of pseudoephedrine is stirred with liquid wax to form a homogeneous mixture, cooled and then forced through a screen to form granules. Preferred matrix materials are waxy substances. Especially preferred are hydrogenated castor oil and carnauba wax and stearyl alcohol.

**Brief Summary Text (84):**

A further preferred process for manufacturing matrix multiparticulates involves using an organic solvent to aid mixing of the pseudoephedrine with the matrix material. This technique can be used when it is desired to utilize a matrix material with an unsuitably high melting point that, if the material were employed in a molten state, would cause decomposition of the drug or of the matrix material, or would result in an unacceptable melt viscosity, thereby preventing mixing of pseudoephedrine with the matrix material. Pseudoephedrine and matrix material may be combined with a modest amount of solvent to form a paste, and then forced through a screen to form granules from which the solvent is then removed. Alternatively, pseudoephedrine and matrix material may be combined with enough solvent to completely dissolve the matrix material, and the resulting solution which may contain solid drug particles spray dried to form the particulate dosage form. This technique is preferred when the matrix material is a high molecular weight synthetic polymer such as a cellulose ether or cellulose ester. Solvents typically employed for the process include acetone, ethanol, isopropanol, ethyl acetate, and mixtures of two or more.

**Brief Summary Text (85):**

Dosage forms according to the invention can be tested to measure the rate of release both of cetirizine and of pseudoephedrine by the following procedure which employs a USP Apparatus 2 as described in USP Chapter <711>. The apparatus is implemented to run at a paddle speed of 50 rpm, with one liter of distilled water at 37.degree. C. The dosage form is added to the apparatus and the cetirizine is monitored by HPLC with UV detection to ascertain when the cetirizine concentration reaches a steady value. Typical times at which the cetirizine concentration is monitored are at 15 minutes, minutes, 45 minutes, and 1 hour after addition, although different or additional times can also be used. The point at which the immediate-release cetirizine concentration reaches a steady value is the time at which release is on the order of 80% complete, typically less than one hour. The pseudoephedrine concentration is similarly monitored at two hour intervals, starting at the first two hour mark after adding the dosage form to the USP Apparatus 2, until the concentration has leveled out.

**Brief Summary Text (86):**

The HPLC system used to monitor the USP aqueous medium can be the same reverse phase system for both cetirizine and pseudoephedrine. The column is typically a C-18 CN column, of the type available under the registered trademark ZORBAX from Mac-Mod. The mobile phase is maintained at 30.degree. C. and can be an isocratic combination of 1:1 (v/v) water/methanol with the pH adjusted to 6.5 by adding sodium dihydrogen phosphate and sodium hydroxide, as appropriate, and made 5 mM in 1-octanesulfonic acid sodium salt. Detection can be effected for both cetirizine and pseudoephedrine with a UV detector at 214 nm. Typical retention times are 3-4 minutes for pseudoephedrine and 8-9 minutes for cetirizine, thus assuring reasonably good separation.

**Detailed Description Text (4):**

A pseudoephedrine tablet core of the composition given below was made by the following procedure. A wet granulation of ingredients 1-5 was prepared with a 50/50 solution of isopropyl alcohol and ethanol, ingredients 6-7. Sufficient solvent addition and mixing of the granulation resulted in a doughy mass that was subsequently crumbled and dried in an oven. The dried granulation was properly sized by comminution. An acceptably sized form of sodium chloride was added to make up 5.6% of the said granulation. To this, 0.5% of magnesium stearate was added to serve as a tableting aid. The granulation was compressed into tablet cores containing approximately 180 mg of pseudoephedrine using 13/32" standard round concave tooling on a conventional rotary tablet press.

**Detailed Description Text (8):**

The dried membrane-coated cores were subsequently coated with the cetirizine HCl-containing coating that also included a 60 mg dose of pseudoephedrine hydrochloride shown in the following table. The drug layer was overcoated onto about 900 grams of membrane coated cores in a perforated tablet coating pan (Hi-Coater, HCT-30, Vector Corp.). The drug solution was sprayed at about 9 g/min while maintaining an outlet temperature of 35-40.degree. C. Enough drug layer was applied so that the tablet cores were overcoated with about 60 mg of pseudoephedrine hydrochloride and about 10 mg of cetirizine hydrochloride.

Detailed Description Text (12):

These overcoated tablets were subjected to the following storage conditions as part of an accelerated stability study (note: "RH" means relative humidity): 50.degree. C./20% RH, 40.degree. C./75% RH and 5.degree. C. After three weeks, the following degradation of cetirizine to the cetirizine ester of glycerol (CEG) was determined.

Detailed Description Text (15):

A pseudoephedrine tablet core of the composition given below was prepared by the following procedure. First, 155 kg of granulation was prepared by combining components 1, 2, and 3 (in the proportions shown) in a 10 ft.sup.3 twin shell blender equipped for wet granulation ("P-K Processor", Patterson-Kelly Corp., East Stroudsburg, Pa. U.S.A.). The blended components were granulated by the addition of water (component 4). Upon completion of water addition, mixing was continued for 3 minutes, and then the wet mass was dried in-situ by application of vacuum at 60.degree. C. with agitation. When the water content of the granulation had reached a level of less than 1% (determined by loss-on-drying measurement), the dried granulation was passed through a mill and divided into two equal portions for subsequent processing. One portion of the dried granulation was then combined with microcrystalline cellulose (component 5) in the P-K Processor, blended for minutes, milled, blended again for 15 minutes and then blended with magnesium stearate (component 6) 5 for minutes to yield the final blend, suitable for tableting on a high speed press. The blend was tableted on a Manesty Mark IIa tablet press using 7/16" standard round concave tooling to yield a batch of tablet cores of nominal weight 535 mg each.

Detailed Description Text (19):

The cores prepared above were further coated with cetirizine by spraying them with the solution shown below in the HiCoater using two spray guns at a solution application rate of 20 g/minute per gun and an outlet air temperature of 45.degree. C. Application of the correct amount of drug to each tablet was verified by measuring the weight gain of a sample of 100 tablets and confirmed by UV analysis. The cetirizine-coated tablets were further coated with the taste-mask coating by spraying them with the taste-mask coating solution (also shown below) at an application rate of 60 g/minute using one spray gun and an outlet air temperature of 45.degree. C. to yield the final product.

Detailed Description Text (20):

Cetirizine Coating Solution

Detailed Description Text (26):

Cetirizine Coating

Detailed Description Text (29):

Pseudoephedrine HCl, microcrystalline cellulose and hydroxypropyl cellulose were blended in a 2 cubic foot P-K processor for minutes, milled through a Fitz mill (using size 2AA plate and knives forward) and blended for minutes. The blend was wet granulated in the P-K processor with 9% water by weight. The granulation was dried in the P-K processor using a jacket temperature of 60.degree. C., to a final water content of less than 1% by loss on drying. The dried granulation was milled through a Fitz mill (using size 2A plate and knives forward), blended for 10 minutes in a 5 cubic foot twin shell blender. Microcrystalline cellulose was added and blended for 15 minutes. Magnesium stearate was added and blended for 5 minutes. The granulation was compressed on a Kilian LX21 tablet press using 7/16" SRC tooling at a tablet hardness of 10 kp. The asymmetric membrane coating was applied to the tablets in an HCT-60 coater using a spray rate of 140 g/min, then dried for 16 hours in a Stokes forced hot air dryer. The asymmetric membrane coated tablets were coated with cetirizine in an HCT-60 coater using one spray nozzle and a spray rate of 40 g/minute. The final taste mask coating was then applied using a spray rate of 40 g/minute. The formulation did not form mono-alcohol esters of cetirizine, since no mono-alcohols reactive to cetirizine were used in the formulation. No significant esters of cetirizine were formed with other hydroxyl-containing excipients.

Detailed Description Text (31):

A pseudoephedrine tablet core of the composition given below was made by the following procedure. Approximately 56 kg of a wet granulation was prepared by first blending ingredients 1-5 in a cubic foot blender for 20 minutes. It was then milled and reblended for another 15 minutes. Half of the blend was transferred to a 140 quart Hobart mixer. The 50/50 solution of isopropyl alcohol and ethanol (about 7.7 kg) was added slowly over 12 minutes such that the blend was thoroughly wetted and actually appeared to be "overwet". The granulation was mixed for about 35 more minutes. The wet mass was tray dried in a forced air oven at 50.degree. C. for 12-16 hours.

Detailed Description Text (36):

The dried tablets were then subsequently coated with the cetirizine HCl-containing coating shown in the following table. The drug layer was overcoated onto about 1100 grams of membrane coated tablets in a perforated tablet coating pan (Hi-Coater, HCT-30, Vector Corp.). The drug solution was sprayed at about 5 g/min while maintaining an outlet temperature of about 40.degree. C. Enough drug layer was applied such that the tablet cores were overcoated with about 10 mg of cetirizine hydrochloride.

Detailed Description Text (40):

These overcoated tablets were subjected to the following storage conditions as part of an accelerated stability study: 50.degree. C., 40.degree. C., 30.degree. C. and 5.degree. C. Degradation to the cetirizine ester of PEG3350 was determined after 12 weeks and 18 months as shown in the table below.

Detailed Description Paragraph Table (1):

% w/w of finished Weight Component Grade blend and tablet (mg/tablet) 1. Pseudoephedrine HCl USP 40.00 180.00 2. Lactose, Anhydrous NF 15.37 69.15 3. Calcium Phosphate Dibasic, Anhydrous USP 24.93 112.20 4. Ethylcellulose, Standard 100 Premium NF 9.07 40.80 5. Povidone (Plasdone C-15) USP 4.53 20.40 6. Isopropyl Alcohol USP (45.00)\* 7. Ethanol, 95% USP (45.00)\* 8. Sodium Chloride USP 5.60 25.20 9. Magnesium Stearate NF 0.50 2.25 Total 100.00 450.00 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (2):

% w/w of coating Weight Component Grade solution (mg/tablet) 1. Pseudoephedrine HCl 180 mg Tablet Core 450.00 2. Ethylcellulose, Standard 10 Premium NF 4.2 81.00 3. Cellulose Acetate NF 0.5 9.00 4. Glycerin 99.5% (Glycerol) USP 2.3 45.00 5. Purified Water USP 2.8 (54.00)\* 6. Isopropyl Alcohol USP 12.4 (239.14)\* 7. Ethanol, 95% USP 22.8 (439.71)\* 8. Acetone NF 55.0 (1060.71)\* Total 100.00 585.00 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (3):

% w/w of Weight Component Grade coating soln (mg/tablet) 1. Pseudoephedrine HCl 180 AM Coated Tablet Pharm 585.00 2. Pseudoephedrine HCl USP 6.5 60.00 3. Cetirizine HCL[] Pharm 1.1 10.00 4. Opadry, Clear (YS-1-7006) Pharm 6.0 55.50 5. Purified Water USP 86.4 (797.30)\* Total 100.00 710.50 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (4):

% w/w of Weight Component Grade coating soln (mg/tablet) 1. Cetirizine HCl/Pseudoephedrine Pharm 710.50 HCl (10/240 mg) Tablet 2. Opadry, Clear (YS-5-1-7006) Pharm 5.0 19.50 3. Purified Water USP 95.0 (370.50)\* Total 100.00 730.00 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (5):

% CEG as determined relative Storage Conditions to the cetirizine peak 50.degree. C./20% RH 57% 40.degree. C./75% RH 43% 5.degree. C. 17%

Detailed Description Paragraph Table (6):

Weight Component (mg/tablet) 1. Pseudoephedrine HCl 240.00 2. Microcrystalline cellulose, NF (Avicel PH 101) 67.48 3. Hydroxypropyl cellulose NF (Klucel EXF) 10.82 4. Purified Water, USP \*(31.80) 5. Microcrystalline cellulose, NF (Avicel PH 200) 213.98 6. Magnesium Stearate, NF 2.67 534.95 mg \*Note: water is used in process, but removed

Detailed Description Paragraph Table (8):

Weight Component (mg/tablet) Cetirizine HCl 10.00 Opadry, Clear (YS-5-19010) 19.30 Purified Water (470.70) 29.3 mg TASTE MASK COATING Weight Component (mg/tablet) Opadry, White (YS-5-18011) 19.70 Purified Water (177.30) 20.20 mg

Detailed Description Paragraph Table (9):

Weight Component (mg/tablet) Pseudoephedrine HCl 240.00 Microcrystalline cellulose, NF (Avicel PH 101) 67.50 Microcrystalline cellulose, NF (Avicel PH 200) 214.00 Hydroxypropyl cellulose, NF 10.82 Purified Water, USP (31.80) Magnesium Stearate, NF 2.67 534.99 mg

Detailed Description Paragraph Table (11):

Weight Component (mg/tablet) Cetirizine HCl 10.00 Opadry, Clear (YS-5-19010) 19.30 Purified Water (303.70) 29.3 mg

Detailed Description Paragraph Table (13):

% w/w of finished Weight Component Grade blend and tablet (mg/tablet) 1. Pseudoephedrine HCl USP 40.00 240.00 2. Lactose, Anhydrous NF 15.37 92.20 3. Calcium Phosphate Dibasic, Anhydrous USP 24.93 149.60 4. Ethylcellulose, Standard



100 Premium NF 9.07 54.40 5. Povidone (Plasdone C-15) USP 4.53 27.20 6. Isopropyl Alcohol USP (90.00)\* 7. Ethanol, 95% USP (90.00)\* 8. Sodium Chloride USP 5.60 33.60 9. Magnesium Stearate NF 0.50 3.00 Total 100.00 600.00 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (14):

% w/w of coating Weight Component Grade solution (mg/tablet) 1. Pseudoephedrine HCl 240 mg Tablet 600.00 Core 2. Ethylcellulose, Standard 100 Premium NF 7.2 76.46 3. Cellulose Acetate NF 0.8 8.50 4. Polyethylene Glycol (Carbowax 3350) NF 3.3 35.04 5. Purified Water USP 3.3 (35.47)\* 6. Isopropyl Alcohol USP 10.6 (112.14)\* 7. Ethanol, 95% USP 22.0 (233.63)\* 8. Acetone NF 52.8 (560.71)\* Total 100.00 720.00 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (15):

% w/w of Weight Component Grade coating soln (mg/tablet) 1. Pseudoephedrine HCl 240 mg Pharm 720.00 AM Coated Tablet 2. Cetirizine HCl Pharm 1.7 10.00 3. Opadry, Clear (YS-5-19010) Pharm 5.8 35.00 4. Purified Water USP 92.5 (555.00)\* Total 100.00 765.00 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (16):

% w/w of Weight Component Grade coating soln (mg/tablet) 1. Cetirizine HCl/Pseudoephedrine Pharm 765.00 HCl (10/240 mg) Tablet 2. Opadry, Clear (YS-5-19010) Pharm 5.0 23.00 3. Purified Water USP 95.0 (437.00)\* Total 100.00 788.00 \*() denotes a volatile component, not present in the final dosage form.

Detailed Description Paragraph Table (17):

% PEG 3350 ester of Cetirizine as determined relative Storage to the cetirizine peak Conditions 12 weeks 18 months 50.degree. C. 0.40 40.degree. C. 0.17 30.degree. C. 0.06 0.09 5.degree. C. ND 0.01 ND = none detected

CLAIMS:

1. A solid dosage form comprising cetirizine and pseudoephedrine

wherein at least a portion of said pseudoephedrine is contained in a core comprising said portion, whereby release of said portion of pseudoephedrine into an environment of use is sustained;

wherein said cetirizine is contained as an immediate-release component in said dosage form; and

wherein said dosage form is substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof, said alcohol and/or derivatives being present in an amount less than that required to react with 5% of the cetirizine in the dosage form.

2. A dosage form as defined in claim 1, wherein said cetirizine is in the form of a pharmaceutically acceptable salt.

4. A dosage form as defined in claim 1, wherein said pseudoephedrine is in the form of a pharmaceutically acceptable salt.

9. A dosage form as defined in claim 1, wherein all of said pseudoephedrine is contained in said core.

10. A dosage form as defined in claim 1, wherein said core comprises a pseudoephedrine-containing immediate release composition surrounded by a rate limiting membrane.

11. A dosage form as defined in claim 1, wherein said core comprises a pseudoephedrine-containing sustained release matrix.

15. A dosage form as defined in claim 1, wherein said portion of pseudoephedrine is released into said environment of use over a period of 4 to 36 hours.

17. A solid dosage form comprising cetirizine and pseudoephedrine, wherein at least a portion of said pseudoephedrine is contained in a core surrounded by a permeable membrane, whereby release of said portion of pseudoephedrine into an environment of use is sustained;

wherein said cetirizine is contained as an immediate-release component in said dosage form; and

wherein said dosage form is substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof, said alcohol and/or derivatives being present in an amount less than that required to react with 5% of the cetirizine in the dosage form.

18. A dosage form as defined in claim 17, wherein said cetirizine is in the form of a pharmaceutically acceptable salt.
20. A dosage form as defined in claim 17, wherein said pseudoephedrine is in the form of a pharmaceutically acceptable salt.
26. A dosage form as defined in claim 25, wherein said core comprises about 60 to about 360 mg of pseudoephedrine, about 70 to about 425 mg of compressible excipients, about 2 to about 17 mg of binder, and about 0.5 to about 4 mg of lubricant.
29. A dosage form as defined in claim 17, wherein said portion of pseudoephedrine is released into said environment of use over a period of 4 to 36 hours.
31. A process for making a solid dosage form comprising cetirizine and pseudoephedrine, comprising coating a sustained release core comprising pseudoephedrine, with an immediate release layer comprising cetirizine and a water soluble film forming polymer, said layer and said core being substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof, said alcohol and/or derivatives being present in an amount less than that required to react with 5% of the cetirizine in the dosage form.
34. A process for making a solid dosage form containing cetirizine and pseudoephedrine, comprising the steps of  
coating a shaped composition comprising pseudoephedrine with a permeable membrane,  
coating said permeable membrane with an immediate release layer comprising cetirizine and a water soluble film forming polymer,  
said composition, said membrane and said layer being substantially free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof.
36. A process as defined in claim 35, wherein said core comprises about 60 to about 360 mg of pseudoephedrine, about 70 to about 425 mg of compressible excipients, about 2 to about 17 mg of binder, and about 0.5 to about 4 mg of lubricant.
39. A dosage form as defined in claim 1, containing an amount of said alcohol and/or derivatives less than the amount required to react with 1% of the cetirizine in the dosage form.
40. A dosage form as defined in claim 39, containing an amount of said alcohol and/or derivatives less than the amount required to react with 0.5% of the cetirizine in the dosage form.
42. A dosage form as defined in claim 17, containing an amount of said alcohol and/or derivatives less than the amount required to react with 1% of the cetirizine in the dosage form.
43. A dosage form as defined in claim 42, containing an amount of said alcohol and/or derivatives less than the amount required to react with 0.5% of the cetirizine in the dosage form.
45. A process as defined in claim 31, wherein said dosage form contains an amount of said alcohol and/or derivatives less than the amount required to react with 1% of the cetirizine therein.
46. A process as defined in claim 45, wherein said dosage form contains an amount of said alcohol and/or derivatives less than the amount required to react with 0.5% of the cetirizine therein.

**WEST**

Generate Collection

Print

L6: Entry 4 of 14

File: USPT

Jan 9, 2001

US-PAT-NO: 6171618

DOCUMENT-IDENTIFIER: US 6171618 B1

TITLE: Combination dosage form comprising cetirizine and pseudoephedrine

DATE-ISSUED: January 9, 2001

## INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Johnson; Barbara A.	Niantic	CT		
Korsmeyer; Richard W.	Old Lyme	CT		
Oksanen; Cynthia A.	Mystic	CT		

## ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE CODE
Pfizer Inc.	New York	NY			02

APPL-NO: 08/ 864490 [PALM]

DATE FILED: May 28, 1997

## PARENT-CASE:

This application is filed claiming priority from co-pending Provisional Application No. 60/015,865, filed May 29, 1996.

INT-CL: [07] A61 K 9/22, A61 K 9/36

US-CL-ISSUED: 424/472; 424/480

US-CL-CURRENT: 424/472; 424/480

FIELD-OF-SEARCH: 424/490, 424/480, 424/474, 424/472

## PRIOR-ART-DISCLOSED:

## U.S. PATENT DOCUMENTS

Search Selected

Search ALL

	PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<input type="checkbox"/>	<u>4792448</u>	December 1988	Ranade	424/438
<input type="checkbox"/>	<u>4915954</u>	April 1990	Ayer et al.	424/473
<input type="checkbox"/>	<u>4996061</u>	February 1991	Webb et al.	424/475
<input type="checkbox"/>	<u>5085865</u>	February 1992	Nayak	424/472
<input type="checkbox"/>	<u>5558879</u>	September 1996	Chen et al.	424/480
<input type="checkbox"/>	<u>5627183</u>	May 1997	Gray	514/255
<input type="checkbox"/>	<u>5654005</u>	August 1997	Chen et al.	424/480
<input type="checkbox"/>	<u>5807579</u>	September 1998	Vikov et al.	424/469

## FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
1220137	April 1987	CA	
1297026	March 1992	CA	
1314485	March 1993	CA	
2147606	May 1994	CA	
2163121	December 1994	CA	
2185893	September 1995	CA	
2217226	October 1996	CA	
2054752	December 1996	CA	
0281708	September 1988	EP	
0294993	December 1988	EP	
0311067	April 1989	EP	
0357369	March 1990	EP	
0396404	November 1990	EP	
0811374A1	October 1997	EP	
0811374	December 1997	EP	
WO 9013295	November 1990	WO	
9202212	February 1992	WO	
9428880	December 1994	WO	
9507103	March 1995	WO	

ART-UNIT: 165

PRIMARY-EXAMINER: Kulkosky; Peter F.

## ABSTRACT:

A dosage form containing cetirizine as an immediate release component and pseudoephedrine or a pharmaceutically acceptable salt thereof as a controlled release component. A portion of the pseudoephedrine can also be incorporated as an immediate release component. The dosage form is free of alcohols having a molecular weight lower than 100 and reactive derivatives thereof.

47 Claims, 0 Drawing figures